


RESEARCH ARTICLE

Accumulation of 3-aminopropylphosphonate in the ex vivo brain observed by phosphorus-31 nuclear magnetic resonance

David Shaul^{1,2} | Benjamin Grieb^{1,3} | Naama Lev-Cohain¹ | Jacob Sosna¹ |
J. Moshe Gomori¹ | Rachel Katz-Brull^{1,2} 

¹Department of Radiology, Hadassah Medical Center, Hebrew University of Jerusalem, The Faculty of Medicine, Jerusalem, Israel

²The Wohl Institute for Translational Medicine, Jerusalem, Israel

³Department of Psychiatry and Psychotherapie I (Weissenau), ZfP Suedwuerttemberg, Ulm University, Ravensburg, Germany

Correspondence

Rachel Katz-Brull, Department of Radiology, Hadassah Medical Center, Hebrew University of Jerusalem, The Faculty of Medicine, Jerusalem, Israel.

Email: rkb@hadassah.org.il

Funding information

Israel Science Foundation; Grant Agreement Number: 1379/18.

European Commission; Grant/Award Number: 858149 - AlternativesToGd.

Jabotinsky Scholarship of the Israeli Ministry of Science and Technology for Applied and Engineering Sciences for Direct PhD Students; Grant/Award Number: 3-15892 (for DS).

3-aminopropylphosphonate (3-APP) is known for its use as an exogenous indicator of extracellular volume and pH in phosphorus-31 nuclear magnetic resonance (³¹P NMR) studies. We used 3-APP for estimating the extracellular volume in NMR studies of several ex vivo preparations including retrograde perfused mouse heart (n = 4), mouse liver slices (n = 2), xenograft breast cancer tumors (n = 7, MCF7), and rat brain slices (n = 4). In the former three preparations, the 3-APP signal was stable in lineshape and intensity for hours and the chemical shift of the signal in the presence of the biological sample was the same as in the perfusion medium without the biological sample. However, in studies of brain slices, the 3-APP signal appeared split into two, with an upfield component (0.7 ± 0.1 ppm to the left) increasing with time and showing a wider linewidth (66.7 ± 12.6 vs. 39.1 ± 7.6 Hz, the latter is of the perfusion medium signal). This finding suggests that 3-APP inadvertently accumulated in brain slices, most likely as a membrane bound form. This observation limits the use of 3-APP as an inert biochemical indicator in brain preparations and should be taken into account when using 3-APP in vivo.

KEYWORDS

3-APP 3-aminopropylphosphonate, ATP adenosine triphosphate, 3-APA 3-aminopropanephosphonic acid, brain, ³¹P phosphorous NMR spectroscopy

1 | INTRODUCTION

3-aminopropylphosphonate (3-APP) is an exogenous marker, used predominantly to measure the extracellular pH of tumors in preclinical NMR studies.¹⁻³ In 1994, it was shown to be nontoxic and membrane impermeant.² Since then, it has been used by numerous groups to measure extracellular pH and to estimate extracellular volume in a variety of animal and tumor models.¹⁻³

We were interested in using 3-APP to characterize extracellular volume in a perfused tissue slices preparation and a perfused mouse heart model that we have been developing for monitoring hyperpolarized substrates metabolism (¹³C-NMR) in tandem with phosphorus-31 nuclear magnetic resonance (³¹P-NMR) monitoring of pH and high energy phosphate status.^{4,5} In such ³¹P-NMR studies, the Pi signal is composed of both the extracellular Pi from the medium perfusing the slices (or heart) and the intracellular Pi within the cells in the slices (or heart). For this reason, it is difficult to estimate the extracellular volume from the Pi signal and there are no other intrinsic markers for the extracellular volume. 3-APP, being membrane impermeant and having a distinguishable chemical shift at a higher field, appeared potentially useful.

Abbreviations used: 3-APA, 3-aminopropanephosphonic acid; 3-APP, 3-aminopropylphosphonate; ATP, adenosine triphosphate.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *NMR in Biomedicine* published by John Wiley & Sons Ltd.

We show here that while 3-APP is useful for determining extracellular space in tissue slices of mice tumors and livers, and in perfused mouse hearts, it is not useful for this purpose in perfused rat cerebrum slices. In the latter, 3-APP accumulated in the slices and for this reason could not serve as a biochemically inert indicator.

2 | METHODS

2.1 | Materials

3-APP was purchased from Sigma-Aldrich (Rehovot, Israel) and was added to the perfusion media of ex vivo preparations at a concentration of 1 mM (heart and brain) and 2 mM (MCF7 breast cancer tumors and liver).

2.2 | Perfused ex vivo preparations

Tissue slices and organs were prepared and perfused in the NMR spectrometer as previously described for mouse liver^{6,7} ($n = 2$), breast cancer xenograft tumors⁸ ($n = 7$), mouse heart⁹ ($n = 4$), and rat cerebrum¹⁰⁻¹² ($n = 4$).

2.3 | Ethics

The joint ethics committee (Institutional Animal Care and Use Committee) of the Hebrew University and Hadassah Medical Center approved the study protocols for animal welfare (protocol numbers: MD-16-14739-1 and MD-19-15882-3 [brain]; MD-17-15045-5 [breast cancer tumors]; MD-16-14741-1 [liver]; and MD-16-14724-1 and MD-19-15827-1 [heart]). The Hebrew University is an AAALAC International accredited institute. Care was taken to minimize pain and discomfort to the animals.

2.4 | NMR spectroscopy

³¹P spectroscopy was performed in a 5.8-T NMR spectrometer equipped with a broadband NMR probe (RS2D, Mundolsheim, France). The tissue was placed inside a 10-mm NMR tube and connected to the perfusion system as previously described.⁶⁻¹² The spectra were acquired using a nutation angle of 50° and a repetition time of 1.1 s. Each spectrum was obtained using 820 scans (total scan time of 15 min).

2.5 | Spectral analysis

Spectral analysis was performed using MNova (Mestrelab Research, Santiago de Compostela, Spain). Each spectrum was zero-filled to 16,384 points, drift-corrected, exponentially multiplied with a 10-Hz line broadening (LB), and baseline corrected. Calculation of the full width at half maximum (FWHM) of the 3-APP signal in the brain slices was performed using the processed spectra (FWHM_{processed}, with 10-Hz LB) because the natural FWHM (FWHM_{natural}) was difficult to assess due to the high noise level in the non line-broadened spectra.

The FWHM_{natural} of these signals was then derived using Equation (1).

$$\text{FWHM}_{\text{natural}} = \text{FWHM}_{\text{processed}} - \text{LB}. \quad (1)$$

We note that because 3-APP accumulation in the brain affected the signal shape and the corresponding linewidth, improvement of signal-to-noise ratio (SNR) by increasing the number of scans would not necessarily provide a better measure of the linewidth.

2.6 | Calculation of 3-APP accumulation in the brain

Deconvolution of the signals of 3-APP in the perfusion medium (3-APP_m) and in the brain (3-APP_b) was performed in Excel (Microsoft, Ra'anana, Israel). The chemical shift of the 3-APP_m signal was the same as its chemical shift in medium that did not contain the slices, and the chemical shift of 3-APP_b was 0.7 ± 0.1 ppm upfield ($n = 4$). In the spectra recorded from the perfused brain slices, a Lorentzian lineshape was centered at the

frequency of the 3-APP_m signal and fitted to its linewidth (39.1 ± 7.6 Hz, $n = 4$). This fitted signal was then subtracted from the total 3-APP signal (3-APP_t) to yield the 3-APP_b signal. Then the integrated intensities of both components were normalized to the integrated intensity of the 3-APP signal recorded from the medium alone (without the brain slices). Because the T_1 of the 3-APP_b signal is unknown, absolute quantification could not be performed.

3 | RESULTS

Ex vivo preparations of brain slices, breast cancer slices (MCF7), liver slices, and a retrograde perfused mouse heart were perfused with media supplemented with 3-APP, as described in the Methods section. ^{31}P NMR spectra of the tissues, recorded under this condition, are shown in Figure 1A (black). The ^{31}P spectra of the perfusion media without the involved tissue are shown in Figure 1A (blue). Figure 1B shows the expanded 3-APP signal for each tissue. In the breast cancer, liver, and heart preparations, the 3-APP signal of the medium alone was greater than the 3-APP signal from the sample that contained the tissues. The difference between these two signals is proportional to the volume that is occupied by the tissue and, by inference, to the proportion of the intracellular volume and medium/extracellular volume detected by the NMR probe.¹ However, in the brain, an additional wider signal (66.7 ± 12.6 Hz, $n = 4$) was observed at a higher field (0.7 ± 0.1 ppm to the left, $n = 4$) relative to the 3-APP signal of the medium.

Figure 2 shows the dynamic nature of the 3-APP signal from perfused rat brain slices. When the perfusion medium was supplemented with 3-APP, a gradual increase in the 3-APP_b signal was observed. When 3-APP was washed out of the perfusion medium, the 3-APP_b signal remained (Figure 2, yellow circle), while the 3-APP_m signal disappeared as expected.

We note that the upfield chemical shift of the 3-APP_b signal is in line with the acidity of the brain tissue in this preparation (approximately 6.9^{pH}), as the 3-APP chemical shift is known to be higher at a lower pH.²

Figure 3A demonstrates the accumulation of 3-APP in brain slices from four different rats (four different experimental days). The part of the ^{31}P spectrum showing this signal is enlarged. The time of 3-APP incubation was 3.2, 4.2, 3.5, and 2.25 h for the slices from animals #1, #2, #3, and #4, respectively (Figure 3A).

Figure 3B shows a typical 3-APP signal deconvolution into its 3-APP_m and 3-APP_b signal components (see also the Methods section). In Figure 3C,D we show the change in these two components over time, in four different brain slice preparations ($n = 4$). It appears that while the 3-APP_m signal remains unchanged over time (Figure 3C), the 3-APP_b signal increases (Figure 3D). During the experimental time frame of about 4.5 h, the accumulation rate did not reach a plateau.

The pH values of the environments in which the 3-APP_m and the Pi signals were present were calculated using the chemical shifts of both signals in the four experiments that were carried out on brain slices. The calculation was performed according to previously described equations.^{1,2} The results are shown in Table 1, and suggest that the two pH determinations are congruent. A slightly lower pH value resulted from the calculation of pH that was based on the chemical shift of the 3-APP_m signal. However, considering the small number of experiments and the SNR of the




FIGURE 1 Phosphorus-31 nuclear magnetic resonance (^{31}P -NMR) spectra acquired from ex vivo preparations of perfused brain slices, breast cancer slices, liver slices, and retrograde perfused heart. (A) Blue: spectra of the perfusion media for the various tissues supplemented with 3-aminopropylphosphonate (3-APP) (without the tissue). Black: spectra of the viable tissues perfused with these media. (B) Expansion of the 3-APP signal of the spectra shown in (A). Only the spectrum of the brain slices shows an additional signal, upfield to the 3-APP signal of the perfusion medium. Each spectrum was recorded using a flip angle of 50° , TR of 1.1 s, and 5740 averages (total scan time of 105 min). Signal assignment: 1, 3-APP; 2, phosphomonoesters, 3, inorganic phosphate; 4, phosphocreatine; 5, γ -adenosine triphosphate (γ -ATP); 6, α -ATP; 7, β -ATP




FIGURE 2 Demonstration of 3-aminopropylphosphonate (3-APP) accumulation in the rat brain slices. Representative ^{31}P spectra of 3-APP signal over time in the course of a wash-in/wash-out experiment. From the bottom up: the medium only (without brain slices) devoid of 3-APP; supplementation of 3-APP to the same medium; time 0 is the time at which the brain slices were introduced to the NMR tube; after 135 min, 3-APP was washed out of the perfusion system by replacing the perfusion medium with a medium that did not contain 3-APP. The yellow straight line indicates the position (in ppm) of the 3-APP_m signal; the yellow circle highlights the 3-APP_b signal (bound to the brain slices). Each spectrum was recorded using a flip angle of 50° , TR of 1.1 s, and 820 averages (total scan time of 15 min)

^{31}P spectra, we hesitate to draw further conclusions from this difference. Therefore, this pH comparison suggested that the two signals represent compounds that are found in the same environment. Based on the medium composition and the signal shape, the Pi signal appears to be predominantly from the perfusion medium. Therefore, it strongly suggests that the 3-APP_m signal is found in the perfusion medium as well, as also suggested by the wash-in/wash-out experiments (Figure 2).

4 | DISCUSSION

Multiple studies have established the use of 3-APP as an indicator of extracellular volume and pH.¹⁻³ In our work, we aimed at using 3-APP to estimate the volume occupied by the tissue cells that is visible to the NMR probe. However, in the perfused rat brain slices preparation, the




FIGURE 3 Accumulation of 3-aminopropylphosphonate (3-APP) in the brain. (A) Accumulation of 3-APP in brain slices from four different rats. For each animal (#1 through 4), the spectrum in black shows the 3-APP_t signal at the beginning of the incubation of the slices with 3-APP, and the signal in blue shows the 3-APP_t at the end of the incubation period. The 3-APP_m and 3-APP_b components of the 3-APP_t signal are denoted as 1 and 2, respectively. The spectra were acquired with 820 scans, within 15 min each. (B) A typical ³¹P signal of 3-APP_t acquired from rat brain slices perfused with a medium containing 3-APP (1 mM). The acquired signal is shown in solid black; a Lorentzian lineshape fitting to the 3-APP_m signal is shown with a dashed line; the result of subtracting 3-APP_m signal from the 3-APP_t signal is shown in gray and comprises the 3-APP_b signal. The spectrum was recorded using a flip angle of 50°, a TR of 1.1 s, and 820 averages (total scan time of 15 min). (C and D) Changes in the 3-APP_m and the 3-APP_b signals, respectively, with perfusion time. The levels of the 3-APP_m and 3-APP_b signals were normalized to the 3-APP signal of the medium alone on each experimental time course (*n* = 4, each experimental time course is marked with a different marker type) recorded at the beginning of each experiment. The trend line in D demonstrates the increase with time of the 3-APP_b signal (*R*² = 0.7)

TABLE 1 Calculated environment pH based on the chemical shifts of the 3-APP_m and Pi signals

Animal number	pH based on the chemical shift of 3-APP _m	pH based on the chemical shift of Pi
1	7.39	7.46
2	7.44	7.50
3	7.39	7.43
4	7.28	7.35
Average ± standard deviation	7.38 ± 0.06*	7.44 ± 0.06*

*Significance was calculated using a two-tailed, paired, Student's *t*-test (*p* = 0.003).

3-APP signal accumulated with time and displayed changes in the signal chemical shift and lineshape. This accumulation did not reach a plateau after 4.5 h of perfusion (Figure 3D). Because 3-APP was not accumulating in the other tissues that we have studied (Figure 1), it is not likely that the accumulation in the perfused brain slices is due to a physical or other barrier that would lead to an increase in the concentration of this compound within the NMR probe. Therefore, we conclude that 3-APP has been accumulating in the brain slices. In addition, the 3-APP_b signal remained constant after 3-APP was removed from the perfusion medium, reinforcing this conclusion (Figure 2). Unfortunately, we could not perform absolute quantification of the rate of 3-APP accumulation in the brain. Bhujwalla et al.¹ reported a longer *T*₁ relaxation for 3-APP in RIF-1 tumors (6.1 s) compared with that of 3-APP in saline (1.7 s). In our experimental setting, evaluation of 3-APP *T*₁ was not possible because the 3-APP_b signal (and likely the content of 3-APP in the brain slices) had increased with time.

The increase in the linewidth of 3-APP_b compared with 3-APP_m suggests that 3-APP_b is membrane bound and not taken up into the intracellular space. This is because NMR linewidths increase with an increase in the molecular weight due to reduced molecular mobility.¹³ The NMR signals of membrane-bound small molecules show the same effect, as they too have longer average rotational correlation times.

We note that the ³¹P signals of phosphodiester were not observed in the rat brain slices. Such signals have been observed in the intact rat^{14,15} and human^{16–19} brain. However, the absence of phosphodiester signals is consistent with earlier studies of rat brain slices.^{20,21} This difference between the in vivo and ex vivo spectra is, to the best of our knowledge, as of yet unexplained.

Previous studies have shown that 3-aminopropanephosphonic acid (3-APA), which is the acidic form of 3-APP, acts as a central and a peripheral GABA_B receptor agonist.²² This binding likely provides an explanation for 3-APP accumulation in the brain. There are two implications of this observation: (1) The accumulation of 3-APP in the brain impairs the ability to correctly evaluate the extracellular volume, as opposed to other tissues that we have experimented with; and (2) GABA_B agonists modulate brain activity and affect cerebral blood flow,^{23–25} thus 3-APP may influence these factors as well, and this should be carefully considered when using 3-APP in vivo.

In another aspect, this observation may open a window for new uses of 3-APP in ³¹P NMR studies of the brain. This accumulation in the brain may be an indicator of pathologies and damaged tissues such as, for example, under stroke. 3-APP may also become a marker of brain hypoxia and cell death due to the sensitivity of its chemical shift to tissue acidity.

In summary, this study demonstrates accumulation of 3-APP in ex vivo perfused rat brain slices, as opposed to other ex vivo perfused tissue preparations (heart, liver, breast cancer tumor). This observation warrants caution as regards its ability to serve as an inert indicator on NMR studies.

ACKNOWLEDGMENTS

The authors wish to thank Drs. Talia Harris, Atara Nardi-Schreiber, Gal Sapir, Yael Adler-Levi, Ayelet Gamliel, and Sivarvanjan Uppala, and Mr. Assad Azar for assistance with data collection in the initial phase of this project. This study was funded by the Israel Science Foundation under grant agreement no. 1379/18, by the European Commission (Grant/Award Number: 858149 - AlternativesToGd), and the Jabotinsky Scholarship of the Israeli Ministry of Science and Technology for Applied and Engineering Sciences for Direct PhD Students (No. 3-15892 for DS).

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Rachel Katz-Brull  <https://orcid.org/0000-0003-4850-1616>

REFERENCES

- Bhujwala ZM, McCoy CL, Glickson JD, Gillies RJ, Stubbs M. Estimations of intra- and extracellular volume and pH by P-31 magnetic resonance spectroscopy: effect of therapy on RIF-1 tumours. *Br J Cancer*. 1998;78(5):606-611. doi:10.1038/bjc.1998.548
- Gillies RJ, Liu Z, Bhujwala Z. P-31 MRS measurements of extracellular pH of tumors using 3-aminopropylphosphonate. *Am J Physiol*. 1994;267(1):C195-C203. doi:10.1152/ajpcell.1994.267.1.C195
- Gillies RJ, Morse DL. In vivo magnetic resonance spectroscopy in cancer. *Annu Rev Biomed Eng*. 2005;7(1):287-326. doi:10.1146/annurev.bioeng.7.060804.100411
- Harris T, Azar A, Sapir G, et al. Real-time ex-vivo measurement of brain metabolism using hyperpolarized 1-C-13 pyruvate. *Sci Rep*. 2018;8(1):9564. doi:10.1038/s41598-018-27747-w
- Harris T, Uppala S, Lev-Cohain N, et al. Hyperpolarized product selective saturating-excitations for determination of changes in metabolic reaction rates in real-time. *NMR Biomed*. 2020;33(2):e4189. doi:10.1002/nbm.4189
- Lev-Cohain N, Sapir G, Harris T, et al. Real-time ALT and LDH activities determined in viable precision-cut mouse liver slices using hyperpolarized 1-C-13 pyruvate-Implications for studies on biopsied liver tissues. *NMR Biomed*. 2019;32(2):e4043. doi:10.1002/nbm.4043
- Lev-Cohain N, Sapir G, Uppala S, et al. Differentiation of heterogeneous mouse liver from HCC by hyperpolarized ¹³C magnetic resonance. *Science*. 2021;3(1):8. doi:10.3390/sci3010008
- Adler-Levy Y, Nardi-Schreiber A, Harris T, et al. In-cell determination of lactate dehydrogenase activity in a luminal breast cancer model - ex vivo investigation of excised xenograft tumor slices using dDNP hyperpolarized 1-C-13 pyruvate. *Sensors*. 2019;19(9):2089. doi:10.3390/s19092089
- Shaul D, Azar A, Sapir G, et al. Correlation between lactate dehydrogenase/pyruvate dehydrogenase activities ratio and tissue pH in the perfused mouse heart: A potential noninvasive indicator of cardiac pH provided by hyperpolarized magnetic resonance. *NMR Biomed*. 2021;34(2):e4444. doi:10.1002/nbm.4444
- Grieb B, Uppala S, Sapir G, Shaul D, Gomori JM, Katz-Brull R. Curbing action potential generation or ATP-synthase leads to a decrease in in-cell pyruvate dehydrogenase activity in rat cerebrum slices. *Sci Rep*. 2021;11(1):10211. doi:10.1038/s41598-021-89534-4
- Sapir G, Shaul D, Lev-Cohain N, Sosna J, Gomori JM, Katz-Brull R. LDH and PDH activities in the ischemic brain and the effect of reperfusion—an ex vivo MR study in rat brain slices using hyperpolarized [1-¹³C]pyruvate. *Metabolites*. 2021;11(4):210. doi:10.3390/metabo11040210
- Shaul D, Grieb B, Sapir G, et al. The metabolic representation of ischemia in rat brain slices: A hyperpolarized ¹³C magnetic resonance study. *NMR Biomed*. 2021;34(7):e4509. doi:10.1002/nbm.4509
- Bradbury JH, Stubbs GJ. Relations between line widths of proton magnetic resonance spectra and molecular weight of poly-γ-benzyl-L-glutamate. *Nature*. 1968;218(5146):1049-1050. doi:10.1038/2181049a0
- Smith CD, Landrum W, Carney JM, Landfield PW, Avison MJ. Brain creatine kinase with aging in F-344 rats: analysis by saturation transfer magnetic resonance spectroscopy. *Neurobiol Aging*. 1997;18(6):617-622. doi:10.1016/S0197-4580(97)00156-5
- Tiret B, Brouillet E, Valette J. Evidence for a "metabolically inactive" inorganic phosphate pool in adenosine triphosphate synthase reaction using localized ³¹P saturation transfer magnetic resonance spectroscopy in the rat brain at 11.7 T. *J Cereb Blood Flow Metab*. 2016;36(9):1513-1518. doi:10.1177/0271678X16657095
- Li S, van der Veen JW, An L, et al. Cerebral phosphoester signals measured by ³¹P magnetic resonance spectroscopy at 3 and 7 Tesla. *PLoS ONE*. 2021;16(3):e0248632. doi:10.1371/journal.pone.0248632
- Peeters TH, van Uden MJ, Rijpmma A, Scheenen TWJ, Heerschap A. 3D ³¹P MR spectroscopic imaging of the human brain at 3 T with a ³¹P receive array: an assessment of ¹H decoupling, T₁ relaxation times, ¹H-³¹P nuclear Overhauser effects and NAD⁺. *NMR Biomed*. 2021;34(5):e4169. doi:10.1002/nbm.4169
- Ren J, Sherry AD, Malloy CR. ³¹P-MRS of healthy human brain: ATP synthesis, metabolite concentrations, pH, and T₁ relaxation times. *NMR Biomed*. 2015;28(11):1455-1462. doi:10.1002/nbm.3384
- Das N, Ren JM, Spence JS, Rackley A, Chapman SB. Relationship of parieto-occipital brain energy phosphate metabolism and cognition using ³¹P MRS at 7-Tesla in amnesic mild cognitive impairment. *Front Aging Neurosci*. 2020;12:222. doi:10.3389/fnagi.2020.00222
- Espanol MT, Litt L, Chang LH, James TL, Weinstein PR, Chan PH. Adult rat brain-slice preparation for nuclear magnetic resonance spectroscopy studies of hypoxia. *Anesthesiology*. 1996;84(1):201-210. doi:10.1097/0000542-199601000-00022

21. Tasker RC, Sahota SK, Williams SR. Bioenergetic recovery following ischemia in brain slices studied by ³¹P-NMR spectroscopy: differential age effect of depolarization mediated by endogenous nitric oxide. *J Cereb Blood Flow Metab.* 1996;16(1):125-133. doi:10.1097/00004647-199601000-00015
22. Ong J, Harrison NL, Hall RG, Barker JL, Johnston GAR, Kerr DIB. 3-aminopropanephosphinic acid is a potent agonist at peripheral and central presynaptic GABA-b receptors. *Brain Res.* 1990;526(1):138-142. doi:10.1016/0006-8993(90)90260-I
23. Franklin TR, Wang Z, Sciortino N, et al. Modulation of resting brain cerebral blood flow by the GABA B agonist, baclofen: A longitudinal perfusion fMRI study. *Drug Alcohol Depend.* 2011;117(2-3):176-183. doi:10.1016/j.drugalcdep.2011.01.015
24. Fergus A, Lee KS. GABAergic regulation of cerebral microvascular tone in the rat. *J Cereb Blood Flow Metab.* 1997;17(9):992-1003. doi:10.1097/00004647-199709000-00009
25. Alborch E, Torregrosa G, Terrasa JC, Estrada C. GABA receptors mediate cerebral vasodilation in the unanesthetized goat. *Brain Res.* 1984;321(1):103-110. doi:10.1016/0006-8993(84)90685-1

How to cite this article: Shaul D, Grieb B, Lev-Cohain N, Sosna J, Gomori JM, Katz-Brull R. Accumulation of 3-aminopropylphosphonate in the ex vivo brain observed by phosphorus-31 nuclear magnetic resonance. *NMR in Biomedicine.* 2022;e4721. doi:10.1002/nbm.4721